

Home Care/ Corporate Office 500 W Lanier Ave Unit 401 Fayetteville, GA 30214 678-962-5850

Inpatient Unit 320 Parkway Dr. NE FL4 Atlanta, GA 30212 (404) 941-1700

Hospice for Liver Disease

Patient has both (A or B):

A. Synthetic failure as demonstrated by i. or ii. and iii.

i. Prothrombin time (PPT) prolonged more than five (5) seconds over control

ii. International Normalized Ration (INR) > 1.5

AND

iii. Serum albumin <2.5gm/dl

AND

B. End -Stage liver disease is present, and the patient has one or more of the following conditions:

- Ascites, refractory to treatment or patient declines or is non-complaint
 History of spontaneous bacterial peritonitis
- Hepatorenal syndrome (elevated creatinine with oliguria [<400 ml/day])
- Hepatic encephalopathy, refractory to treatment or patient non-complaint
 History of recurrent variceal bleeding despite intensive therapy or patient declines therapy

Supporting evidence for hospice eligibility:

- Progressive malnutrition
- · Muscle wasting with reduced strength
- Ongoing alcoholism (>80 gm ethanol/day)
 - Hepatocellular carcinoma
- Hepatitis B surface antigen positive
- Hepatitis C refractory to interferon treatment.

In the absence of one or more of these findings, rapid decline or comorbidities may also support eligibility for hospice care.

Harbor Grace Hospice | www.harborgracehospice.com







